


Long term variability of Cen X–3 with MAXI

J. J. Rodes, T. Mihara, J. M. Torrejón, S. Nakahira, G. Bernabéu, G. Sanjurjo



Universitat d'Alacant
Universidad de Alicante



Abstract

Cen X–3 is an O-type donor Supergiant X-ray Binary with a neutron star as a compact object. The aim of this work is to study both the light curve and orbital phase spectroscopy of this system in the long term. Here we estimate the **orbital period from the light curve** and then we derive the good time interval to extract the spectrum from different orbital phases. We obtain **orbital phase-averaged** and **phase-resolved spectra** and we analyse the variability of the model parameters to infer properties of the circumstellar environment of Cen X-3 and possible dependences of the spectral parameters. The MAXI spectra in the 2-20 keV energy range were fitted with **a model of Comptonisation of cool photons on hot electrons modified by a partial covering absorption and a fluorescence iron emission line at 6.4 keV**.


Introduction

The eclipsing high mass X-ray binary (HMXB) pulsar Cen X-3 has an O-type supergiant companion, V779 Cen. The orbital period of the binary system is ~ 2.1 days with a low eccentricity of ≤ 0.0016 [1] and the magnetised neutron star has a spin period of ~ 4.8 s [7]. The distance to the source was estimated to be ~ 8 kpc [4], although [9] gave a value of 5.7 ± 1.5 kpc.


The **Monitor of All Sky X-ray Image** (MAXI) presents both all-sky coverage and moderate energy resolution, which gives us the possibility to investigate the orbital light curves and the orbital phased-resolved spectra of **Cen X–3** [6]. MAXI observing strategy suppress effectively the short term variability associated with accretion and enhances the long term, permanent structures present in the stellar wind and circumsource environment [2], [3].

Timing analysis

First, we have obtained the MAXI/GSC on-demand light-curve of Cen X–3 from MJD 55 058 to MJD 57 315 (i.e. more than 6 years) in the 2–20 keV energy band. Then, we have searched for a **period** assuming a sinusoidal signal and derived a $P_{\text{orb}} = 2.0870 \pm 0.0006$ **days** which is consistent with the value given by [5].




a) The extraction region




b) Background subtracted light curves

Spectral analysis

We have extracted the orbital phase-averaged spectrum of Cen X–3 with MAXI/GSC using the MAXI on-demand processing, carefully excluding any contamination by nearby brighter sources. We tested both phenomenological and physical models commonly applied to accreting X-ray pulsars. We rebinned all extracted spectra to obtain spectral bins Gaussian distributed. The best fit was attained using COMPST, in XSPEC terminology [8], modified by a partial covering model implementation of the Tübingen-Boulder absorption model (*TBnew*) which includes the most up to date absorption cross sections and abundances [10]. We needed to put an additional Gaussian absorption line at ~ 5 keV to compensate the incompleteness of the GSC response (Nakahira private communication). Moreover, the fluorescence iron emission line at ~ 6.4 keV was also present in the spectrum.




a) Averaged spectrum



b) Phase-resolved spectra

Then we have obtained **orbital phase-resolved spectra** of the HMXB pulsar Cen X–3 accumulating the 60 s duration scans into 10 orbital phase bins. We have fitted the orbital phase-resolved spectra with the same model we used in the orbital phase averaged spectrum. Nevertheless, no absorption Gaussian component at 5 keV was needed in these fits. We note that the optical counterpart blocked most of the X-ray emission in eclipse in all the energy bands, so there is no scattering in the wind. For the **fluorescence iron emission line**, a Gaussian profile is assumed. Moreover, this iron line is **not detected** both in the **eclipse and egress spectra** nor resolved the complex iron line in the energy range 6.4–6.7 keV.

Orbital phase-resolved spectra



The **unabsorbed flux** appears nearly constant in the orbital phase range $[0.1 - 0.4]$, $F_{(2-20) \text{ keV}} \sim 4.4 \times 10^9 \text{ erg cm}^{-2} \text{ s}^{-1}$, showing that **the accretion rate is quite stable**. The relative flux during the eclipse is a factor of 4 smaller than during out of eclipse.

The **equivalent width** (EW) of the Fe K α is consistent with a constant value of $\sim 220 \pm 60 \text{ eV}$ taking into account the uncertainties. The centre of the energy of the iron line changes with orbital phase suggesting the **coexistence of two iron lines at different energies**.

References

- [1] Bildsten, L. et al. 1997 *ApJS*, 113, 367
- [2] Doroshenko, V. et al. 2013 *A&A*, 554, A37
- [3] Islam, N. & Paul B. 2014 *MNRAS*, 441, 2539
- [4] Krzeminski, W. 1974 *ApJ*, 192, L135
- [5] Nagase, F., Corbet, R. H. D. et al. 1992 *ApJ*, 396, 147
- [6] Rodes-Roca, J. J., Mihara, T. et al. 2016 *A&A*, to be submitted
- [7] Schreier, E., Levinson, R. et al. 1972 *ApJ*, 172, L79
- [8] Sunyaev, R. A. & Titarchuk, L. G. 1980 *A&A*, 86, 121
- [9] Thompson, T. W. J. & Rothschild, R. E. 2009 *ApJ*, 691, 1744
- [10] Wilms, J., Allen, A. & McCray, R. 2000 *ApJ*, 542, 914

Funding

This work was supported by the project ESP2014-53672-C3-3-P and partially by GV2014/088. TM acknowledges the support by the grant by the University of Alicante under visiting research programme INV15-08. JJRR acknowledges the support by the Matsumae International Foundation Research Fellowship No14G04.